

NOTAS DE MATEMATICA

Nº 45

A NOTE ON SPECTRAL AND PRESPECTRAL OPERATORS

BY

T.V. PANCHAPAGESAN

DEPARTAMENTO DE MATEMATICA

FACULTAD DE CIENCIAS

UNIVERSIDAD DE LOS ANDES

MERIDA - VENEZUELA

1981

# A NOTE ON SPECTRAL AND PRESPECTRAL OPERATORS

BY

T.V. PANCHAPAGESAN<sup>+</sup>

ABSTRACT. Results known for spectral and prespectral operators in weakly complete Banach spaces are extended to Banach spaces with Bessaga-Pelczyński property.

1. PRELIMINARIES. In this note by a Banach space we mean a non-zero complex Banach space. Because of varying terminology in literature, we give the following definition.

DEFINITION 1.1. Let  $\Sigma$  be a  $\sigma$ -algebra of subsets of a non-empty set  $S$  and  $X$  a Banach space. If  $\Gamma$  is a total set of  $X^*$ , the dual of  $X$ , and  $E(\cdot)$  is a Boolean homomorphism of  $\Sigma$  on a Boolean algebra (B.A.) of projections on  $X$ , we say that  $E(\cdot)$  is a spectral measure of class  $(\Sigma, \Gamma)$  when  $E(\Sigma)$  is norm bounded in the Banach algebra  $B(X)$  of all bounded operators in  $X$  and when  $x^* E(\cdot) x$  is countably additive on  $\Sigma$ , for each  $x \in X$  and  $x^* \in \Gamma$ .

---

<sup>+</sup>Supported by C.D.C.H. projects C-80-149,150 of the Universidad de los Andes, Mérida, Venezuela.

AMS subject classification-1979-47 B40-spectral operators, decomposable operators etc.

When  $E(\cdot)$  is a spectral measure of class  $(\Sigma, X^*)$ , by Orlicz-Pettis' theorem,  $E(\cdot)$  is countably additive on  $\Sigma$  in the strong operator topology and in this case  $E(\cdot)$  is called a spectral measure on  $\Sigma$ .

An operator  $T \in B(X)$  is called a prespectral operator of class  $\Gamma$  if there is a spectral measure  $E(\cdot)$  of class  $(\Sigma_p, \Gamma)$  where  $\Gamma$  is total in  $X^*$  and  $\Sigma_p$  is the  $\sigma$ -algebra of all Borel sets in  $\mathbb{C}$ , such that for each  $\delta$  in  $\Sigma_p$ ,  $E(\delta)T = TE(\delta)$  and the spectrum of  $T$  restricted to  $E(\delta)X$  is contained in the closure of  $\delta$ . Then  $E(\cdot)$  is called a resolution of the identity of class  $\Gamma$  for  $T$ .

A prespectral operator  $T$  of class  $X^*$  is called a spectral operator. In this case,  $T$  has a unique resolution of the identity.

DEFINITION 1.2. A Banach space  $X$  is said to have Bessaga-Pelczyński (B-P) property if every weakly unconditionally convergent series in  $X$  is strongly unconditionally convergent.

LEMMA 1.3. Let  $\Omega$  be a compact  $T_2$  space and  $X$  a Banach space with B-P property. Then every bounded linear transformation  $T: C(\Omega) \rightarrow X$  is weakly compact, where  $C(\Omega)$  is the Banach algebra of all complex continuous functions on  $\Omega$ .

PROOF. By theorem 5 of Bessaga and Pelczyński [1], the hypothesis on  $X$  is equivalent to the fact that  $X$  does not contain

an isomorphic copy of  $c_0$ . Now the lemma follows from Theorem 15, p. 160 of Diestel and Uhl [2].

2. MAIN RESULTS. In what follows  $X$  will denote a complex Banach space with B-P property. Since a weakly complete Banach space has B-P property (See [1]) all the results in this section hold for weakly complete Banach spaces also.

THEOREM 2.1. Let  $A$  be an algebra of operators in  $X$  and let it be the image under a continuous homomorphism  $\phi$  of the Banach algebra  $C(\Omega)$  of all complex functions on a compact Hausdorff space  $\Omega$ . Then there exists a uniquely determined spectral measure  $E(\cdot)$  in  $X$  on the Borel sets of  $\Omega$ , for which

$$\phi(f) = \int_{\Omega} f(\lambda) E(d\lambda), \quad f \in C(\Omega).$$

PROOF. For a fixed  $x \in X$ , consider the map  $T_x: C(\Omega) \rightarrow X$ , defined by  $T_x f = \phi(f)x$ . Since  $T_x$  is a continuous linear transformation and  $X$  has B-P property, by Lemma 1.3 we have that  $T_x$  is weakly compact. Now following the same argument as in the proof of Theorem 2.5 of Chapter XVII of Dunford and Schwartz [4], the present theorem is established.

We use the terms 'complete' and ' $\sigma$ -complete' B.A of projections on  $X$  in the sense of Definition 3.1 of Chapter XVII of Dunford and Schwartz [4].

LEMMA 2.2. A strongly closed bounded Boolean algebra of projections in  $X$  is complete.

PROOF. The proof of Lemma 3.5 of Chapter XVII of Dunford and Schwartz [4] holds here, except that we appeal to Theorem 2.1. above, in stead of Theorem 2.5 of Chapter XVII of [4].

THEOREM 2.3. A bounded B.A. of projections on  $X$  is complete if and only if it is strongly closed.

PROOF. The theorem follows from Lemma 2.2. above and from Corollary 3.7 of Chapter XVII of [4].

COROLLARY 2.4. Every bounded B.A. of projections  $B$  on  $X$  can be embedded in a  $\sigma$ -complete B.A. of projections on  $X$ , contained in  $\bar{B}^s$  (the closure in the strong operator topology).

PROOF. When  $B$  is norm bounded by a finite positive number  $M$ , then it is easily verified that  $\bar{B}^s$  is a B.A. of projections and that  $\|E\| \leq M, E \in \bar{B}^s$ . Now the corollary follows from Theorem 2.3.

The above corollary leads to the following interesting result about prespectral operators.

THEOREM 2.5. Every prespectral operator  $T$  on  $X$  is a spectral operator.

PROOF. Let  $T$  be a prespectral operator of class  $\Gamma$  on  $X$ ,

with  $E(\cdot)$  a resolution of the identity of class  $\Gamma$ . Then by Definition 1.1 and by Corollary 2.4 there exists a  $\sigma$ -complete B.A.  $B$  of projections on  $X$ , which contains the range of  $E(\cdot)$  on  $\Sigma_p$ . If  $\{\delta_n\}$  is a sequence of mutually disjoint sets in  $\Sigma_p$ , then  $F_n = \sum_{k=1}^n E(\delta_k)$  is in  $B$  and  $\{F_n\}$  is a non-decreasing sequence of projections in  $B$ . Therefore, by Lemma 3.4 of Chapter XVII of [4], for each  $x \in X$

$$\lim_n F_n x = \left( \bigvee_1^\infty F_n \right) x .$$

In other words,

$$(1) \quad \left( \bigvee_1^\infty F_n \right) x = \lim_{n \rightarrow \infty} \sum_1^n E(\delta_k) x, \quad x \in X.$$

But,  $x^* E(\cdot) x$  is countably additive for each  $x \in X$  and  $x^* \in \Gamma$  and therefore from (1), we have

$$(2) \quad x^* \left( \bigvee_1^\infty F_n \right) x = \sum_{k=1}^\infty x^* E(\delta_k) x = x^* E \left( \bigcup_1^\infty \delta_k \right) x$$

$x \in X, x^* \in \Gamma.$

Because  $\Gamma$  is total, (1) and (2) give that

$$(3) \quad \left( \bigvee_1^\infty F_n \right) x = E \left( \bigcup_1^\infty \delta_k \right) x, \quad x \in X .$$

From (1) and (3) it follows that  $E(\cdot)$  is countably additive in the strong operator topology. Hence  $T$  is spectral.

THEOREM 2.6. Let  $T_1$  and  $T_2$  be two commuting spectral operators on  $X$ , with  $E_1(\cdot)$  and  $E_2(\cdot)$  as their respective resolutions of the identity and  $S_1$  and  $S_2$  as their respective scalar parts. If the B.A. of projections  $B$  determined by  $E_1(\cdot)$  and  $E_2(\cdot)$  is bounded, then  $T_1 + T_2$  and  $T_1 T_2$  are spectral with  $S_1 + S_2$  and  $S_1 S_2$  as their respective scalar parts and  $G_1(\cdot)$  and  $G_2(\cdot)$  as their respective resolutions of the identity, where

$$G_1(\delta)x = \int E_2(\delta-\lambda)E_1(d\lambda)x,$$

$$G_2(\delta)x = \int E_2(\delta/\lambda)E_1(d\lambda)x$$

for each  $x \in X$  and  $\delta \in \Sigma_p$ .

PROOF. Let  $T_1 = S_1 + N_1$  and  $T_2 = S_2 + N_2$  be the canonical decompositions of  $T_1$  and  $T_2$ . Then by the generalized Fuglede's theorem (Corollary 3.7, Chapter XV of [4])  $S_1, S_2, N_1$  and  $N_2$  commute with each other and hence  $N_1 + N_2$  and  $N_1 S_2 + N_2 S_1 + N_1 N_2$  are quasi-nilpotent. Now the theorem follows from Theorem 4.5 of Chapter XV of [4], Corollary 2.4 above and Theorem 10 of Panchapagesan [5].

REMARK 1. In view of Theorems 2.1 and 2.3 and Corollary 2.4, Theorems 6.10, 6.12, 6.14, 6.21 and 6.22 and note 6.15 of Dowson [3] and Corollaries 2.12, 2.13 and 2.14 and Theorems 3.18, 3.19 and 3.20 and Corollary 3.28 of Chapter XVII

of Dunford and Schwartz [4], which are known to be true for weakly complete Banach spaces, continue to be valid for Banach spaces with B-P property.

It is also possible to extend the theorems of characterizations of spectral operators in weakly complete Banach spaces to operators in Banach spaces with B-P property. To this end we need the following theorem.

THEOREM 2.7. Let  $E(\cdot): R \rightarrow B(X)$  be a Boolean homomorphism of the algebra of sets  $R$  on a B.A. of projections on  $X$  and let  $E(\cdot)$  be countably additive in the strong operator topology on  $R$ , in the sense that whenever  $\{\delta_n\}$  is a pairwise disjoint sequence of members of  $R$  with  $\bigcup_1^\infty \delta_n \in R$ , then

$$E\left(\bigcup_1^\infty \delta_n\right)x = \sum_1^\infty E(\delta_n)x, \quad x \in X.$$

Then a necessary and sufficient condition that  $E(\cdot)$  can be extended uniquely to a spectral measure  $\bar{E}(\cdot)$  on  $S(R)$ , the  $\sigma$ -algebra generated by  $R$ , is that the range of  $E(\cdot)$  is norm bounded in  $B(X)$ .

PROOF. If such an extension  $\bar{E}(\cdot)$  on  $S(R)$  exists, then as in the second part of the proof of Corollary 3.10 of Chapter XVII of [4] we see that  $\bar{E}(S(R))$  is a  $\sigma$ -complete B.A. of projections on  $X$  and hence is norm bounded by Lemma 3.3 of Chapter XVII of [4]. Conversely, if  $E(R)$  is norm bounded in



$B(X)$ , then  $E(R)$  is a bounded B.A. of projections on  $X$ . Hence from Theorem 8 of Panchapagesan [5] and Corollary 2.4 above, the sufficiency of the condition follows.

REMARK 2. As a consequence of the above Theorem, Theorems 3.14, 4.5, 5.2 and 5.15 of Chapter XVI of Dunford and Schwartz [4], which are known true for weakly complete Banach spaces, continue to be valid for Banach spaces with B-P property.

## R E F E R E N C E S

1. C. Bessaga and A. Pelczyński, On bases and unconditional convergence of series in Banach spaces, *Studia Math.* 17 (1958), 151-164.
2. J. Diestel and J.J. Uhl, *Vector measures*, Amer. Math. Soc. Survery 15 (1977).
3. H.R. Dowson, *Spectral theory of linear operators*, Academic Press, New York, (1978).
4. N. Dunford and J.T. Schwartz, *Linear operators, Part III, Spectral operators*, Interscience, New York (1971).
5. T.V. Panchapagesan, Extension of spectral measures, *Illinois J. Math.* 16 (1972), 130-142.

DEPARTAMENTO DE MATEMATICA  
FACULTAD DE CIENCIAS  
UNIVERSIDAD DE LOS ANDES  
MERIDA, VENEZUELA